

MAKING HEALTHY HAPPEN™
Shaklee

Shaklee®

Omega-3 Gellys™

STRAWBERRY LEMONADE



EPA + DHA chewable Gellys for the whole family.
Great-tasting, sugar free, and readily absorbed.
Promotes brain, heart, eye, joint, immune, and skin health.*

30
GELLYS

300 mg
EPA+DHA PER GELLY

Dr. Shaklee

DIETARY SUPPLEMENT

NET WT. 1.6 OZ. (45 g)
0.05 OZ. (1.5 g) EACH

XD843A

Item #21403

Give Your Body What Every Body Needs

At every age and every stage, the body requires omega-3s to operate optimally. This potent formula contains pure EPA + DHA to help support:

- Developing & Aging Brains*
 - Heart Health*
 - Eye Health*
 - Joint Health*
 - Healthy Immune Systems*
 - Skin Health*
- + **Sustainably sourced and no fishy aftertaste**

Superior Delivery with 3x More DHA + EPA†

Developed by Norwegian scientists, this revolutionary technology uses emulsified droplets to mirror the distribution of nutrients in food, leading to improved digestion and absorption of omega-3s.*

*Per Gelly compared to other leading Gummies



NON
GMO

GF
GLUTEN
FREE

SOY
FREE

DAIRY
FREE

ZERO
SUGAR

NO Artificial Flavors or Preservatives

Supplement Facts

Serving Size: 1 or 2 Gellys
Servings Per Container: 30

	Kids (4-18)		Adults (18+)	
	1 Gelly	% DV	2 Gellys	% DV
Calories	5		10	
Total Fat	0.5 g	<1%**	1 g	1%**
Total Omega-3 Fatty Acids	335 mg	†	670 mg	†
DHA (Docosahexaenoic Acid)	250 mg	†	500 mg	†
EPA (Eicosapentaenoic Acid)	50 mg	†	100 mg	†

**Percent Daily Values (DV) are based on a 2,000 calorie diet.
†Daily Value not established.

Directions: Kids take 1 daily with food. Adults take 2 daily with food.

Other Ingredients: Fish Oil Concentrate, Xylitol, Purified Water, Sorbitol, Gelatin, Natural Flavors, Trisodium Citrate, Citric Acid, Beta-Carotene for color.

Distributed by Shaklee Corporation
4747 Willow Rd, Pleasanton, CA 94588
Product questions: 925.734.3638

Manufactured in a facility that may also process tree nuts, milk, egg, soy, wheat, shellfish, and fish.

CAUTION: If taking blood-thinning medications, consult your physician prior to use. Discontinue use two weeks before surgery.

†THESE STATEMENTS HAVE NOT BEEN EVALUATED BY THE FOOD AND DRUG ADMINISTRATION. THIS PRODUCT IS NOT INTENDED TO DIAGNOSE, TREAT, CURE, OR PREVENT ANY DISEASE.